

**Appl. No.** : 09/988,728  
**Filed** : November 16, 2001

## **AMENDMENTS TO THE CLAIMS**

1. (Currently Amended) A method of conducting an assay, the method comprising:

providing a sample of cells in a chamber in a disc, the chamber including at least one capture zone with a capture agent, the disc including at least one inlet port and a vent port on a first surface of the disc;

loading the disc into an optical reader;

rotating the disc so as to separate different cell types into different capture zones;

directing an incident beam of electromagnetic radiation to the capture zone;

detecting a beam of electromagnetic radiation formed after interacting with the disc at the capture zone;

converting the detected beam into an output signal; and

analyzing the output signal to extract therefrom information relating to the number of cells captured at the capture zone

counting captured cells in each of the capture zones; and

providing an output including the counts, wherein the output includes counts for CD4 cells and CD8 cells.

2. (Previously Presented) The method according to claim 1, wherein the chamber is internal to the disc and is bounded on opposite sides by a substrate and cap.

3. (Original) The method according to claim 1, wherein the optical disc is constructed with a reflective layer such that light directed to the capture zone and not striking a cell is reflected.

4. (Previously Presented) The method according to claim 1, wherein the optical disc is constructed such that light directed to the capture zone and not striking a cell is transmitted through the optical disc, the disc being between the light source and ~~the~~ a detector.

5. (Previously Presented) The method according to any one of claims 1-5, wherein the disc surface is coated with a first group of cell capture agents.

**Appl. No.** : **09/988,728**  
**Filed** : **November 16, 2001**

6. (Previously Presented) The method according to claim 5, wherein the cell capture agents define a capture zone.
7. (Previously Presented) The method according to claim 6, wherein a second group of cell capture agents define a second capture .
8. (Original) The method according to claim 7, wherein the first and second captures zones are in one chamber.
9. (Original) The method according to claim 5, wherein the cell capture agents are for binding with cell surface antigen.
10. (Original) The method according to claim 9, wherein the cell surface antigen is selected from the CD family of antigens.
11. (Original) The method according to claim 10, wherein the cell surface antigen is selected from the group consisting of CD3, CD4, CD8, and CD45.
12. (Original) The method according to claim 1, further including:  
directing the sample of cells into proximity with the cell capture agents;  
incubating the cells in the presence of the capture agents; and  
allowing the cells to specifically bind to the capture agents.
13. (Original) The method according to claim 12, further including analyzing the number of cells captured to thereby determine a cell concentration in the sample.
14. (Previously Presented) The method of claim 13, wherein the analyzing includes detecting sufficiently large changes in a level of light reflected from or transmitted through the disc.
15. (Original) The method of claim 13, wherein the analyzing includes using image recognition to count captured cells.

**Appl. No.** : **09/988,728**  
**Filed** : **November 16, 2001**

16. (Previously Presented) The method of claim 15, wherein the image recognition distinguishes one type of white blood cell from another.

17. (Original) The method of claim 1, wherein the chamber has a plurality of capture zones, each having a different cell capture agent.

18. (Previously Presented) The method of claim 17, wherein the rotating includes rotating for a sufficient period of time at a sufficient speed so that the cells have an opportunity to bind with capture molecules.

19. (Original) The method of claim 18, wherein the rotating includes rotating for a sufficient period of time at a sufficient speed so that unbound cells are moved away from the capture zones.

20. (Original) The method of claim 19, wherein the rotating is done at a single speed.

21. (Original) The method of claim 17, further comprising counting ~~the~~ captured cells in each of the capture zones and providing an output including the counts.

22. (Previously Presented) The method of claim 21, wherein the output includes a ratio of CD4 to CD8 cells.

23-29. Cancelled.

30. (Previously Presented) The method of claim 12, wherein the analyzing includes detecting sufficiently large changes in the level of light reflected from or transmitted through the disc.

31. (Previously Presented) The method of claim 12, wherein the analyzing includes using image recognition to count captured cells.

**Appl. No.** : **09/988,728**  
**Filed** : **November 16, 2001**

32. (New) The method of Claim 1, wherein the disc comprises a first layer of streptavidin, a first antibody raised in a first species against a type of immunoglobulin of a second species, and a second antibody raised in the second species against a cell surface antigen.

33. (New) The method of Claim 1, wherein each of the capture zones are sequentially located in a fluid path between the inlet port and the vent port, and wherein capture zones are sequentially provided for CD4, CD8 and a control in relation to the fluid path.